

BAŞARIR M., ŞAHİN A.

**ON THE STRONG AND Δ -CONVERGENCE FOR TOTAL ASYMPTOTICALLY
NONEXPANSIVE MAPPINGS ON A $CAT(0)$ SPACE**

In this paper we give the strong and Δ -convergence theorems of the modified S-iteration and the modified two-step iteration processes for total asymptotically nonexpansive mappings on a $CAT(0)$ space. Our results extend and improve the corresponding recent results announced by many authors in the literature.

Key words and phrases: $CAT(0)$ space, total asymptotically nonexpansive mapping, strong convergence, Δ -convergence, iterative process, fixed point.

Department of Mathematics, Faculty of Sciences and Arts, Sakarya University, Esentepe Campus, 54187, Serdivan, Sakarya, Turkey

E-mail: basarir@sakarya.edu.tr (Başarır M.), ayuce@sakarya.edu.tr (Şahin A.)

INTRODUCTION

A metric space X is a $CAT(0)$ space if it is geodesically connected and every geodesic triangle in X is at least as “thin” as its comparison triangle in the Euclidean plane. Fixed point theory in a $CAT(0)$ space has been first studied by Kirk (see [9, 10]). He showed that every nonexpansive mapping defined on a bounded closed convex subset of a complete $CAT(0)$ space always has a fixed point. Since then the fixed point theory for various mappings in a $CAT(0)$ space has been rapidly developed and a lot of papers have appeared (see [4, 5, 6, 17, 18]).

Nanjaras and Panyanak [13] proved the demiclosedness principle for asymptotically nonexpansive mappings and gave the Δ -convergence theorem of the modified Mann iteration process for mappings of this type in a $CAT(0)$ space. Recently, Chang et. al. [3] introduced total asymptotically nonexpansive mappings and proved the demiclosedness principle for mappings of this type in a $CAT(0)$ space. Also, they presented the Δ -convergence theorem of the modified Mann iteration process for total asymptotically nonexpansive mappings in a $CAT(0)$ space.

In this paper, motivated by the above results, we get some results which are related to the strong and Δ -convergence of the modified S-iteration and the modified two-step iteration processes for total asymptotically nonexpansive mappings on a $CAT(0)$ space. Our results extend and improve the corresponding ones announced by Chang et. al. [3], Khan and Abbas [8], Nanjaras and Panyanak [13] and many others.

YDK 519.6

2010 Mathematics Subject Classification: 47H09, 47H10, 54H25, 58C30.

1 PRELIMINARIES AND LEMMAS

Let (X, d) be a metric space, K be a nonempty subset of X and let $T : K \rightarrow K$ be a mapping. Recall that T is said to be a nonexpansive mapping if

$$d(Tx, Ty) \leq d(x, y), \quad \forall x, y \in K.$$

The map T is said to be an asymptotically nonexpansive mapping if there exists a sequence $\{k_n\} \subset [1, \infty)$ with $k_n \rightarrow 1$ such that

$$d(T^n x, T^n y) \leq k_n d(x, y), \quad \forall n \in \mathbb{N}, x, y \in K.$$

The map T is said to be a uniformly L -Lipschitzian mapping if there exists a constant $L > 0$ such that

$$d(T^n x, T^n y) \leq L d(x, y), \quad \forall n \in \mathbb{N}, x, y \in K.$$

Chang et. al. [3] defined the concept of total asymptotically nonexpansive mapping as follows.

Definition 1 ([3, Definition 2.1]). *Let (X, d) be a metric space, K be a nonempty subset of X and let $T : K \rightarrow K$ be a mapping. T is said to be a total asymptotically nonexpansive mapping if there exist non-negative real sequences $\{\mu_n\}, \{v_n\}$ with $\mu_n \rightarrow 0, v_n \rightarrow 0$ and a strictly increasing continuous function $\zeta : [0, \infty) \rightarrow [0, \infty)$ with $\zeta(0) = 0$ such that*

$$d(T^n x, T^n y) \leq d(x, y) + v_n \zeta(d(x, y)) + \mu_n$$

for all $n \in \mathbb{N}$ and $x, y \in K$.

Remark 1. *From the definitions, it is clear that each nonexpansive mapping is an asymptotically nonexpansive mapping with $k_n = 1, \forall n \in \mathbb{N}$, each asymptotically nonexpansive mapping is a total asymptotically nonexpansive mapping with $\mu_n = 0, v_n = k_n - 1, \forall n \in \mathbb{N}$, $\zeta(t) = t, t \geq 0$, and each asymptotically nonexpansive mapping is a uniformly L -Lipschitzian mapping with $L = \sup_{n \in \mathbb{N}} \{k_n\}$.*

We now give the definition and collect some basic properties of the $CAT(0)$ space.

Let (X, d) be a metric space. A *geodesic path* joining $x \in X$ and $y \in X$ (or more briefly, a *geodesic* from x to y) is a map $c : [0, l] \subset \mathbb{R} \rightarrow X$ such that $c(0) = x, c(l) = y$ and $d(c(t), c(t')) = |t - t'|$ for all $t, t' \in [0, l]$. In particular, c is an isometry and $d(x, y) = l$. The image of c is called a *geodesic* (or *metric*) *segment* joining x and y . When it is unique, this geodesic is denoted by $[x, y]$. The space (X, d) is said to be a *geodesic space* if every two points of X are joined by a geodesic, and X is said to be a *uniquely geodesic space* if there is exactly one geodesic joining x and y for all $x, y \in X$.

A *geodesic triangle* $\Delta(x_1, x_2, x_3)$ in a geodesic metric space (X, d) consist of three points in X (the vertices of Δ) and three geodesic segments joining each pair of vertices (the edges of Δ). A *comparison triangle* for the geodesic triangle $\Delta(x_1, x_2, x_3)$ in (X, d) is a triangle $\bar{\Delta}(x_1, x_2, x_3) = \Delta(\bar{x}_1, \bar{x}_2, \bar{x}_3)$ in the Euclidean plane \mathbb{R}^2 such that $d_{\mathbb{R}^2}(\bar{x}_i, \bar{x}_j) = d(x_i, x_j)$ for $i, j \in \{1, 2, 3\}$. Such a triangle always exists (see [2]).

A geodesic space is said to be a $CAT(0)$ space [2] if all geodesic triangles of appropriate size satisfy the following comparison axiom.

$CAT(0)$: Let Δ be a geodesic triangle in X and let $\bar{\Delta}$ be a comparison triangle for Δ . Then, Δ is said to satisfy the $CAT(0)$ inequality if for all $x, y \in \Delta$ and all comparison points $\bar{x}, \bar{y} \in \bar{\Delta}$,

$$d(x, y) \leq d_{\mathbb{R}^2}(\bar{x}, \bar{y}).$$

Let $x, y \in X$, and by Lemma 2.1(iv) of [6] for each $t \in [0, 1]$, there exists a unique point $z \in [x, y]$ such that

$$d(x, z) = td(x, y), \quad d(y, z) = (1 - t)d(x, y). \quad (1)$$

From now on, we will use the notation $(1 - t)x \oplus ty$ for the unique point z satisfying (1). By using this notation, Dhompongsa and Panyanak [6] obtained the following lemma which will be used frequently in the proof of our main results.

Lemma 1 ([6, Lemma 2.4]). *Let X be a $CAT(0)$ space. Then*

$$d((1 - t)x \oplus ty, z) \leq (1 - t)d(x, z) + td(y, z)$$

for all $t \in [0, 1]$ and $x, y, z \in X$.

In 1976 Lim [12] introduced the concept of Δ -convergence in a general metric space. In 2008 Kirk and Panyanak [11] specialized Lim's concept to the $CAT(0)$ space and proved that it is very similar to the weak convergence in a Banach space. Also, Dhompongsa and Panyanak [6] obtained the Δ -convergence theorems for the Picard, Mann and Ishikawa iterations in a $CAT(0)$ space for nonexpansive mappings.

Let $\{x_n\}$ be a bounded sequence in a $CAT(0)$ space X . For $x \in X$, we set $r(x, \{x_n\}) = \limsup_{n \rightarrow \infty} d(x, x_n)$. The asymptotic radius $r(\{x_n\})$ of $\{x_n\}$ is given by

$$r(\{x_n\}) = \inf\{r(x, \{x_n\}) : x \in X\}$$

and the asymptotic center $A(\{x_n\})$ of $\{x_n\}$ is the set

$$A(\{x_n\}) = \{x \in X : r(x, \{x_n\}) = r(\{x_n\})\}.$$

It is known that in a complete $CAT(0)$ space, $A(\{x_n\})$ consists of exactly one point (see [5, Proposition 7]).

Definition 2 ([11, 12]). *A sequence $\{x_n\}$ in a $CAT(0)$ space X is said to be Δ -convergent to $x \in X$ if x is the unique asymptotic center of $\{u_n\}$ for every subsequence $\{u_n\}$ of $\{x_n\}$. In this case we write $\Delta\text{-}\lim_{n \rightarrow \infty} x_n = x$ and x is called the Δ -limit of $\{x_n\}$.*

Lemma 2. *i) Every bounded sequence in a complete $CAT(0)$ space always has a Δ -convergent subsequence (see [11, p. 3690]).*

ii) Let K be a nonempty closed convex subset of a complete $CAT(0)$ space and let $\{x_n\}$ be a bounded sequence in K . Then the asymptotic center of $\{x_n\}$ is in K (see [4, Proposition 2.1]).

Lemma 3 ([6, Lemma 2.8]). *If $\{x_n\}$ is a bounded sequence in a complete $CAT(0)$ space with $A(\{x_n\}) = \{x\}$, $\{u_n\}$ is a subsequence of $\{x_n\}$ with $A(\{u_n\}) = \{u\}$ and the sequence $\{d(x_n, u)\}$ converges, then $x = u$.*

In [3] it is proved demiclosedness principle for total asymptotically nonexpansive mappings in a $CAT(0)$ space as follows.

Lemma 4 ([3, Theorem 2.8]). *Let K be a closed convex subset of a complete $CAT(0)$ space X and let $T : K \rightarrow K$ be a total asymptotically nonexpansive and uniformly L -Lipschitzian mapping. Let $\{x_n\}$ be a bounded sequence in K such that $\lim_{n \rightarrow \infty} d(x_n, Tx_n) = 0$ and $\Delta\text{-}\lim_{n \rightarrow \infty} x_n = p$. Then $Tp = p$.*

The following lemma is crucial in the study of iteration processes in metric spaces.

Lemma 5 ([14, Lemma 2]). *Let $\{a_n\}$, $\{b_n\}$ and $\{\delta_n\}$ be sequences of non-negative real numbers satisfying the inequality*

$$a_{n+1} \leq (1 + \delta_n)a_n + b_n.$$

If $\sum_{n=1}^{\infty} \delta_n < \infty$ and $\sum_{n=1}^{\infty} b_n < \infty$, then $\lim_{n \rightarrow \infty} a_n$ exists.

Lemma 6 ([13, Lemma 4.5]). *Let X be a $CAT(0)$ space, $x \in X$ be a given point and let $\{t_n\}$ be a sequence in $[b, c]$ with $b, c \in (0, 1)$ and $0 < b(1 - c) \leq \frac{1}{2}$. Let $\{x_n\}$ and $\{y_n\}$ be any sequences in X such that*

$$\limsup_{n \rightarrow \infty} d(x_n, x) \leq r, \quad \limsup_{n \rightarrow \infty} d(y_n, x) \leq r, \quad \lim_{n \rightarrow \infty} d((1 - t_n)x_n \oplus t_n y_n, x) = r$$

for some $r \geq 0$. Then

$$\lim_{n \rightarrow \infty} d(x_n, y_n) = 0.$$

Agarwal, O'Regan and Sahu [1] introduced the modified S-iteration process which is independent of those of the modified Mann iteration [16] and the modified Ishikawa iteration [19]. We apply this iteration process into a $CAT(0)$ space as

$$\begin{cases} x_1 \in K \\ x_{n+1} = (1 - a_n)T^n x_n \oplus a_n T^n y_n \\ y_n = (1 - b_n)x_n \oplus b_n T^n x_n, \quad n \in \mathbb{N}. \end{cases} \tag{2}$$

By taking $T^n = T$ for all $n \in \mathbb{N}$ in (2), we obtain the S-iteration process which is introduced in [1].

Thianwan [20] introduced the two-step iteration process in a Banach space. We give the modified two-step iteration process in a $CAT(0)$ space as follows

$$\begin{cases} x_1 \in K \\ x_{n+1} = (1 - a_n)y_n \oplus a_n T^n y_n \\ y_n = (1 - b_n)x_n \oplus b_n T^n x_n, \quad n \in \mathbb{N}. \end{cases} \tag{3}$$

If $b_n = 0$ for each $n \in \mathbb{N}$, then (3) reduces to the modified Mann iteration process. By taking $T^n = T$ for all $n \in \mathbb{N}$ in (3), we obtain the two-step iteration process.

Our purpose in this paper is to get some results on the strong and Δ -convergence of the modified S-iteration and the modified two-step iteration processes for total asymptotically non-expansive mappings in a $CAT(0)$ space. It is worth mentioning that our results in a $CAT(0)$ space can be applied to any $CAT(k)$ space with $k \leq 0$ since any $CAT(k)$ space is a $CAT(m)$ space for every $m \geq k$ (see [2, p. 165]).

2 MAIN RESULTS

We will denote the set of fixed points of T by $F(T)$, that is, $F(T) = \{x \in X : Tx = x\}$. Firstly, we prove the Δ -convergence theorem of the modified S-iteration process in a $CAT(0)$ space.

Theorem 1. *Let K be a nonempty bounded closed convex subset of a complete $CAT(0)$ space X , $T : K \rightarrow K$ be a total asymptotically nonexpansive and uniformly L -Lipschitzian mapping with $F(T) \neq \emptyset$ and let $\{x_n\}$ be a sequence defined by (2). If the following conditions are satisfied:*

$$(i) \sum_{n=1}^{\infty} v_n < \infty, \sum_{n=1}^{\infty} \mu_n < \infty, \sum_{n=1}^{\infty} a_n < \infty;$$

(ii) *there exists a constant $M^* > 0$ such that $\zeta(r) \leq M^*r, r \geq 0$;*

(iii) *$\{b_n\}$ is the sequence in $[0, 1]$;*

(iv) $\sum_{n=1}^{\infty} \sup \{d(z, T^n z) : z \in B\} < \infty$ *for each bounded subset B of K ;*

(v) *there exist constants $b, c \in (0, 1)$ with $0 < b(1 - c) \leq \frac{1}{2}$ such that $\{a_n\} \subset [b, c]$,*

then the sequence $\{x_n\}$ Δ -converges to a fixed point of T .

Proof. We divide the proof of Theorem 1 into three steps.

Step I. First we prove that for each $p \in F(T)$ the following limit $\lim_{n \rightarrow \infty} d(x_n, p)$ exists. In fact for each $p \in F(T)$, by Lemma 1, we have

$$\begin{aligned} d(y_n, p) &= d((1 - b_n)x_n \oplus b_n T^n x_n, p) \leq (1 - b_n)d(x_n, p) + b_n d(T^n x_n, p) \\ &\leq (1 - b_n)d(x_n, p) + b_n \{d(x_n, p) + v_n \zeta(d(x_n, p)) + \mu_n\} \\ &\leq (1 + b_n v_n M^*) d(x_n, p) + b_n \mu_n \leq (1 + v_n M^*) d(x_n, p) + \mu_n. \end{aligned}$$

Also, we obtain

$$\begin{aligned} d(x_{n+1}, p) &= d((1 - a_n)T^n x_n \oplus a_n T^n y_n, p) \leq (1 - a_n)d(T^n x_n, p) + a_n d(T^n y_n, p) \\ &\leq (1 - a_n) \{d(x_n, p) + v_n \zeta(d(x_n, p)) + \mu_n\} + a_n L d(y_n, p) \\ &\leq (1 - a_n) \{(1 + v_n M^*) d(x_n, p) + \mu_n\} + a_n L \{(1 + v_n M^*) d(x_n, p) + \mu_n\} \\ &= \{(1 - a_n)(1 + v_n M^*) + a_n L(1 + v_n M^*)\} d(x_n, p) + (1 + a_n(L - 1))\mu_n \\ &= \{1 + a_n(L - 1) + v_n M^*(1 + a_n(L - 1))\} d(x_n, p) + (1 + a_n(L - 1))\mu_n. \end{aligned}$$

It follows from condition (i) and Lemma 5 that $\lim_{n \rightarrow \infty} d(x_n, p)$ exists.

Step II. Next we prove that

$$\lim_{n \rightarrow \infty} d(x_n, Tx_n) = 0. \quad (4)$$

In fact, it follows from Step I that for all $p \in F(T)$, $\lim_{n \rightarrow \infty} d(x_n, p)$ exists, so we can assume that

$\lim_{n \rightarrow \infty} d(x_n, p) = r$. Since

$$\begin{aligned} d(T^n y_n, p) &= d(T^n y_n, T^n p) \leq d(y_n, p) + v_n \zeta(d(y_n, p)) + \mu_n \leq (1 + v_n M^*) d(y_n, p) + \mu_n \\ &\leq (1 + v_n M^*) \{(1 + v_n M^*) d(x_n, p) + \mu_n\} + \mu_n \\ &= (1 + v_n M^*) (1 + v_n M^*) d(x_n, p) + (2 + v_n M^*) \mu_n, \end{aligned}$$

we have $\limsup_{n \rightarrow \infty} d(T^n y_n, p) \leq r$. Similarly, we obtain $\limsup_{n \rightarrow \infty} d(T^n x_n, p) \leq r$. On the other hand, since $\lim_{n \rightarrow \infty} d((1 - a_n)T^n x_n \oplus a_n T^n y_n, p) = \lim_{n \rightarrow \infty} d(x_{n+1}, p) = r$, by Lemma 6, we have

$$\lim_{n \rightarrow \infty} d(T^n x_n, T^n y_n) = 0. \quad (5)$$

Since $d(x_{n+1}, T^n x_n) \leq d((1 - a_n)T^n x_n \oplus a_n T^n y_n, T^n x_n) \leq a_n d(T^n y_n, T^n x_n)$ from (5), we obtain

$$\lim_{n \rightarrow \infty} d(x_{n+1}, T^n x_n) = 0. \quad (6)$$

From condition (iv), we have

$$\lim_{n \rightarrow \infty} d(x_n, T^n x_n) = 0. \quad (7)$$

Hence from (6) and (7) we get

$$\lim_{n \rightarrow \infty} d(x_n, x_{n+1}) = 0. \quad (8)$$

Since T is a uniformly L -Lipschitzian mapping, from (7) and (8) we have that

$$\begin{aligned} d(x_n, Tx_n) &\leq d(x_n, x_{n+1}) + d(x_{n+1}, T^{n+1}x_{n+1}) + d(T^{n+1}x_{n+1}, T^{n+1}x_n) + d(T^{n+1}x_n, Tx_n) \\ &\leq d(x_n, x_{n+1}) + d(x_{n+1}, T^{n+1}x_{n+1}) + Ld(x_{n+1}, x_n) + Ld(T^n x_n, x_n) \\ &= (1 + L)d(x_n, x_{n+1}) + d(x_{n+1}, T^{n+1}x_{n+1}) + Ld(T^n x_n, x_n) \rightarrow 0 \quad (\text{as } n \rightarrow \infty). \end{aligned}$$

The equation (4) is proved.

Step III. To show that the sequence $\{x_n\}$ Δ -converges to a fixed point of T , we prove that

$$W_\Delta(x_n) = \bigcup_{\{u_n\} \subset \{x_n\}} A(\{u_n\}) \subseteq F(T)$$

and $W_\Delta(x_n)$ consists of exactly one point. Let $u \in W_\Delta(x_n)$. Then there exists a subsequence $\{u_n\}$ of $\{x_n\}$ such that $A(\{u_n\}) = \{u\}$. By Lemma 2, there exists a subsequence $\{v_n\}$ of $\{u_n\}$ such that $\Delta\text{-}\lim_{n \rightarrow \infty} v_n = v \in K$. By Lemma 4, $v \in F(T)$. Since $\{d(u_n, v)\}$ converges, by Lemma 3, $u = v$. This shows that $W_\Delta(x_n) \subseteq F(T)$. Now we prove that $W_\Delta(x_n)$ consists of exactly one point. Let $\{u_n\}$ be a subsequence of $\{x_n\}$ with $A(\{u_n\}) = \{u\}$ and let $A(\{x_n\}) = \{x\}$. We have already seen that $u = v$ and $v \in F(T)$. Finally, since $\{d(x_n, v)\}$ converges, by Lemma 3, $x = v \in F(T)$. This shows that $W_\Delta(x_n) = \{x\}$. \square

Now we give an example of such mappings which are total asymptotically nonexpansive and uniformly L -Lipschitzian as in Theorem 1.

Let \mathbb{R} be the real line with the usual norm $|\cdot|$ and let $K = [-1, 1]$. Define two mappings $T, S : K \rightarrow K$ by

$$T(x) = \begin{cases} -2 \sin \frac{x}{2}, & \text{if } x \in [0, 1] \\ 2 \sin \frac{x}{2}, & \text{if } x \in [-1, 0) \end{cases} \quad \text{and} \quad S(x) = \begin{cases} x, & \text{if } x \in [0, 1] \\ -x, & \text{if } x \in [-1, 0). \end{cases}$$

It is proved in [7, Example 3.1] that both T and S are asymptotically nonexpansive mappings. Therefore they are total asymptotically nonexpansive and uniformly L -Lipschitzian mappings. Additionally, $F(T) = \{0\}$ and $F(S) = \{x \in K; 0 \leq x \leq 1\}$.

We give the characterization of strong convergence for the modified S-iteration process on a $CAT(0)$ space as follows.

Theorem 2. Let $X, K, T, \{a_n\}, \{b_n\}, \{x_n\}$ satisfy the hypotheses of Theorem 1. Then the sequence $\{x_n\}$ converges strongly to a fixed point of T if and only if

$$\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0,$$

where $d(x, F(T)) = \inf \{d(x, p) : p \in F(T)\}$.

Proof. Necessity is obvious. Conversely, suppose that $\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$. As proved in Theorem 1 (Step I), for all $p \in F(T)$,

$$d(x_{n+1}, p) \leq \{1 + a_n(L - 1) + v_n M^*(1 + a_n(L - 1))\} d(x_n, p) + (1 + a_n(L - 1))\mu_n.$$

This implies that

$$d(x_{n+1}, F(T)) \leq \{1 + a_n(L - 1) + v_n M^*(1 + a_n(L - 1))\} d(x_n, F(T)) + (1 + a_n(L - 1))\mu_n.$$

By Lemma 5, $\lim_{n \rightarrow \infty} d(x_n, F(T))$ exists. Thus by hypothesis $\lim_{n \rightarrow \infty} d(x_n, F(T)) = 0$.

Next, we show that $\{x_n\}$ is a Cauchy sequence in K . Let $\varepsilon > 0$ be arbitrarily chosen. Since $\lim_{n \rightarrow \infty} d(x_n, F(T)) = 0$, there exists a positive integer n_0 such that

$$d(x_n, F(T)) < \frac{\varepsilon}{4}$$

for all $n \geq n_0$. In particular, $\inf \{d(x_{n_0}, p) : p \in F(T)\} < \frac{\varepsilon}{4}$. Thus, there exists $p^* \in F(T)$ such that

$$d(x_{n_0}, p^*) < \frac{\varepsilon}{2}.$$

Now, for all $m, n \geq n_0$, we have

$$d(x_{n+m}, x_n) \leq d(x_{n+m}, p^*) + d(x_n, p^*) \leq 2d(x_{n_0}, p^*) < 2\left(\frac{\varepsilon}{2}\right) = \varepsilon.$$

Hence $\{x_n\}$ is a Cauchy sequence in the closed subset K of a complete $CAT(0)$ space and so it must be convergent to a point q in K . Now, $\lim_{n \rightarrow \infty} d(x_n, F(T)) = 0$ gives that $d(q, F(T)) = 0$ and closedness of $F(T)$ forces q to be in $F(T)$. This completes the proof. \square

Senter and Dotson [15] introduced the concept of *Condition (I)* as follows.

Definition 3 ([15, p. 375]). A mapping $T : K \rightarrow K$ is said to satisfy *Condition (I)* if there exists a non-decreasing function $f : [0, \infty) \rightarrow [0, \infty)$ with $f(0) = 0$ and $f(r) > 0$ for all $r > 0$ such that

$$d(x, Tx) \geq f(d(x, F(T))), \quad \forall x \in K.$$

With respect to the above definition, we have the following strong convergence theorem.

Theorem 3. Let $X, K, T, \{a_n\}, \{b_n\}, \{x_n\}$ satisfy the hypotheses of Theorem 1 and let T be a mapping satisfying *Condition (I)*. Then the sequence $\{x_n\}$ converges strongly to a fixed point of T .

Proof. As proved in Theorem 2, $\lim_{n \rightarrow \infty} d(x_n, F(T))$ exists. Also, by Theorem 1 (Step II), we have $\lim_{n \rightarrow \infty} d(x_n, Tx_n) = 0$. It follows from Condition (I) that

$$\lim_{n \rightarrow \infty} f(d(x_n, F(T))) \leq \lim_{n \rightarrow \infty} d(x_n, Tx_n) = 0.$$

That is, $\lim_{n \rightarrow \infty} f(d(x_n, F(T))) = 0$. Since $f : [0, \infty) \rightarrow [0, \infty)$ is a non-decreasing function satisfying $f(0) = 0$ and $f(r) > 0$ for all $r > 0$, we obtain

$$\lim_{n \rightarrow \infty} d(x_n, F(T)) = 0.$$

The conclusion now follows from Theorem 2. \square

Remark 2. Theorems 1–3 contain some results of Khan and Abbas [8, Theorems 1–3] since each nonexpansive mapping is a total asymptotically nonexpansive mapping.

Now, we give the Δ -convergence theorem of the modified two-step iteration process in a $CAT(0)$ space.

Theorem 4. Let $X, K, T, \{b_n\}$ satisfy the hypotheses of Theorem 1, $\{a_n\}$ be a sequence in $[0, 1]$ and let $\{x_n\}$ be a sequence defined by (3). If the conditions (i)–(iv) in Theorem 1 are satisfied, then the sequence $\{x_n\}$ Δ -converges to a fixed point of T .

Proof. First we prove that for all $p \in F(T)$ the following limit $\lim_{n \rightarrow \infty} d(x_n, p)$ exists. As proved in Theorem 1, we have

$$d(y_n, p) \leq (1 + v_n M^*) d(x_n, p) + \mu_n. \quad (9)$$

Since T is a uniformly L -Lipschitzian mapping, from (9) we have

$$\begin{aligned} d(x_{n+1}, p) &= d((1 - a_n)y_n \oplus a_n T^n y_n, p) \\ &\leq (1 - a_n)d(y_n, p) + a_n d(T^n y_n, p) \\ &\leq (1 - a_n)d(y_n, p) + a_n L d(y_n, p) \\ &= (1 + a_n(L - 1)) d(y_n, p) \\ &\leq (1 + a_n(L - 1)) \{(1 + v_n M^*) d(x_n, p) + \mu_n\} \\ &= \{1 + a_n(L - 1) + v_n M^*(1 + a_n(L - 1))\} d(x_n, p) + (1 + a_n(L - 1))\mu_n. \end{aligned}$$

It follows from Lemma 5 that $\lim_{n \rightarrow \infty} d(x_n, p)$ exists. Next we prove that $\lim_{n \rightarrow \infty} d(x_n, Tx_n) = 0$. From condition (iv), we have

$$\lim_{n \rightarrow \infty} d(x_n, T^n x_n) = \lim_{n \rightarrow \infty} d(y_n, T^n y_n) = 0. \quad (10)$$

By the above equality, we get

$$d(T^n x_n, T^n y_n) \leq L d(x_n, y_n) \leq L b_n d(x_n, T^n x_n) \leq L d(x_n, T^n x_n) \rightarrow 0 \quad (\text{as } n \rightarrow \infty). \quad (11)$$

Since

$$d(x_{n+1}, T^n y_n) \leq d((1 - a_n)y_n \oplus a_n T^n y_n, T^n y_n) \leq (1 - a_n)d(y_n, T^n y_n)$$

from (10), we obtain

$$\lim_{n \rightarrow \infty} d(x_{n+1}, T^n y_n) = 0. \quad (12)$$

From (10), (11) and (12) we have that

$$d(x_n, x_{n+1}) \leq d(x_n, T^n x_n) + d(T^n x_n, T^n y_n) + d(T^n y_n, x_{n+1}) \rightarrow 0 \quad (\text{as } n \rightarrow \infty).$$

The rest of the proof follows the pattern of the Theorem 1 and is therefore omitted. \square

Remark 3. *Theorem 4 contains the main result of Chang et. al. [3, Theorem 3.5] since the modified two-step iteration reduces to the modified Mann iteration. Also, Theorem 4 contains the main result of Nanjaras and Panyanak [13, Theorem 5.7] since each asymptotically nonexpansive mapping is a total asymptotically nonexpansive mapping.*

Finally, we give following theorems related to the strong convergence of the modified two-step iteration process which their proofs are similar arguments of Theorem 2 and Theorem 3, respectively.

Theorem 5. *Let $X, K, T, \{a_n\}, \{b_n\}, \{x_n\}$ satisfy the hypotheses of Theorem 4. Then the sequence $\{x_n\}$ converges strongly to a fixed point of T if and only if $\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$.*

Theorem 6. *Let $X, K, T, \{a_n\}, \{b_n\}, \{x_n\}$ satisfy the hypotheses of Theorem 4 and let T be a mapping satisfying Condition (I). Then the sequence $\{x_n\}$ converges strongly to a fixed point of T .*

REFERENCES

- [1] Agarwal R.P., O'Regan D., Sahu D.R. *Iterative construction of fixed points of nearly asymptotically nonexpansive mappings*. J. Nonlinear Convex Anal. 2007, **8** (1), 61–79.
- [2] Bridson M., Haefliger A. *Metric spaces of non-positive curvature*. Springer-Verlag, Berlin, 1999.
- [3] Chang S.S., Wang L., Joesph Lee H.W., Chan C.K., Yang L. *Demiclosed principle and Δ -convergence theorems for total asymptotically nonexpansive mappings in $CAT(0)$ spaces*. Appl. Math. Comput. 2012, **219** (5), 2611–2617. doi:10.1016/j.amc.2012.08.095
- [4] Dhompongsa S., Kirk W.A., Panyanak B. *Nonexpansive set-valued mappings in metric and Banach spaces*. J. Nonlinear Convex Anal. 2007, **8** (1), 35–45.
- [5] Dhompongsa S., Kirk W.A., Sims B. *Fixed point of uniformly lipschitzian mappings*. Nonlinear Anal.: Theory, Methods & Appl. 2006, **65** (4), 762–772. doi:10.1016/j.na.2005.09.044
- [6] Dhompongsa S., Panyanak B. *On Δ -convergence theorems in $CAT(0)$ spaces*. Comput. & Math. Appl. 2008, **56** (10), 2572–2579. doi:10.1016/j.camwa.2008.05.036
- [7] Guo W.P., Cho Y.J., Guo W. *Convergence theorems for mixed type asymptotically nonexpansive mappings*. Fixed Point Theory Appl. 2012, **2012**:224. doi:10.1186/1687-1812-2012-224
- [8] Khan S.H., Abbas M. *Strong and Δ -convergence of some iterative schemes in $CAT(0)$ spaces*. Comput. & Math. Appl. 2011, **61** (1), 109–116. doi:10.1016/j.camwa.2010.10.037
- [9] Kirk W.A. *Geodesic geometry and fixed point theory*. In: Seminar of Mathematical Analysis (Malaga/Seville, 2002/2003), Colec. Abierta, Univ. Sevilla Secr. Publ., Seville 2003, **64**, 195–225.
- [10] Kirk W.A. *Geodesic geometry and fixed point theory II*. In: International Conference on Fixed Point Theory Appl., Yokohama Publ., Yokohama, 2004, 113–142.
- [11] Kirk W.A., Panyanak B. *A concept of convergence in geodesic spaces*. Nonlinear Anal.: Theory, Methods & Appl. 2008, **68** (12), 3689–3696. doi:10.1016/j.na.2007.04.011
- [12] Lim T.C. *Remarks on some fixed point theorems*. Proc. Amer. Math. Soc. 1976, **60**, 179–182. doi:10.1090/S0002-9939-1976-0423139-X
- [13] Nanjaras B., Panyanak B. *Demiclosed principle for asymptotically nonexpansive mappings in $CAT(0)$ spaces*. Fixed Point Theory Appl. 2010, **2010**:268780. doi:10.1155/2010/268780
- [14] Qihou L. *Iterative sequences for asymptotically quasi-nonexpansive mappings with error member*. J. Math. Anal. Appl. 2001, **259** (1), 18–24. doi:10.1006/jmaa.2000.7353

- [15] Senter H.F., Dotson W.G. *Approximating fixed points of nonexpansive mappings*. Proc. Amer. Math. Soc. 1974, **44** (2), 375–380. doi:10.1090/S0002-9939-1974-0346608-8
- [16] Schu J. *Weak and strong convergence to fixed points of asymptotically nonexpansive mappings*. Bull. Austral. Math. Soc. 1991, **43** (1), 153–159. doi:10.1017/S0004972700028884
- [17] Şahin A., Başarır M. *On the strong and Δ -convergence theorems for nonself mappings on a $CAT(0)$ space*. In: Proc. of the 10th IC-FPTA, Cluj-Napoca, Romania, July 9–18, 2012, 227–240.
- [18] Şahin A., Başarır M. *On the strong convergence of a modified S -iteration process for asymptotically quasi-nonexpansive mappings in a $CAT(0)$ space*. Fixed Point Theory Appl. 2013, **2013**:12. doi:10.1186/1687-1812-2013-12
- [19] Tan K.K., Xu H.K. *Fixed point iteration processes for asymptotically nonexpansive mappings*. Proc. Amer. Math. Soc. 1994, **122** (3), 733–739. doi:10.1090/S0002-9939-1994-1203993-5
- [20] Thianwan S. *Common fixed points of new iterations for two asymptotically nonexpansive nonself-mappings in a Banach space*. J. Comput. Appl. Math. 2009, **224** (2), 688–695. doi:10.1016/j.cam.2008.05.051

Received 20.05.2013

Revised 04.07.2013

Башарір М., Шагін А. *Про сильну і Δ -збіжність для тотальних асимптотично нерозширюваних відображень на $CAT(0)$ простір // Карпатські математичні публікації. — 2013. — Т.5, №2. — С. 170–179.*

В цій статті ми доводимо теореми про сильну і Δ -збіжність модифікованих S -ітерацій і модифікованих двокрокових ітераційних процесів для тотальних асимптотично нерозширюваних відображень на $CAT(0)$ простір. Наші результати розширюють і покращують відповідні недавні результати, що анонсовані багатьма авторами в літературі.

Ключові слова і фрази: $CAT(0)$ простір, тотальне асимптотично нерозширюване відображення, сильна збіжність, Δ -збіжність, ітераційний процес, нерухома точка.

Башарир М., Шагин А. *О сильной и Δ -сходимости для тотальных асимптотически нерасширяющихся отображений на $CAT(0)$ пространство // Карпатские математические публикации. — 2013. — Т.5, №2. — С. 170–179.*

В этой статье мы доказываем теоремы о сильной и Δ -сходимости модифицированных S -итераций и модифицированных двухшаговых итерационных процессов для тотальных асимптотически нерасширяющихся отображений на $CAT(0)$ пространство. Наши результаты расширяют и улучшают соответствующие недавние результаты, анонсированные многими авторами в литературе.

Ключевые слова и фразы: $CAT(0)$ пространство, тотальное асимптотически нерасширяющееся отображение, сильная сходимость, Δ -сходимость, итерационный процесс, неподвижная точка.