



On the analytic extension of the Horn's hypergeometric function H_4

Dmytryshyn R.^{1,✉}, Lutsiv I.-A.¹, Dmytryshyn M.²

The paper establishes new convergence domains of branched continued fraction expansions of Horn's hypergeometric function H_4 with real and complex parameters. These domains enabled the PC method to establish the analytical extension of analytical functions to their expansions in the studied domains of convergence. A few examples are provided at the end to illustrate this.

Key words and phrases: branched continued fraction, Horn hypergeometric function, approximation by rational functions, convergence, analytic continuation.

¹ Vasyl Stefanyk Precarpathian National University, 57 Shevchenka str., 76018, Ivano-Frankivsk, Ukraine

² West Ukrainian National University, 11 Lvivska str., 46009, Ternopil, Ukraine

✉ Corresponding author

E-mail: dmytryshynr@hotmail.com (Dmytryshyn R.), lutsiv.ilona@gmail.com (Lutsiv I.-A.), m.dmytryshyn@wunu.edu.ua (Dmytryshyn M.)

Introduction

The analytic theory of branched continued fractions provides a useful means for the representation and continuation of special functions of one and several variables (see [3, 5, 6, 22–24, 30, 33]), used in mathematical physics. Many applications of branched continued fractions were made in various areas of numerical analysis and theoretical physics, chemistry and engineering [26–28, 34]. This paper continues the works [4, 18–20] and establishes new domains of the analytical continuation of Horn's hypergeometric function H_4 with certain conditions on real and complex parameters, using their representation in branched continued fractions.

Note that the Horn's hypergeometric function H_4 is defined as (see [25])

$$H_4(a, b; c, d; \mathbf{z}) = \sum_{r,s=0}^{\infty} \frac{(a)_{2r+s} (b)_s z_1^r z_2^s}{(c)_r (d)_s r! s!}, \quad |z_1| < p, |z_2| < q, \quad (1)$$

where $a, b, c, d \in \mathbb{C}$, $c, d \notin \{0, -1, -2, \dots\}$, $(\alpha)_0 = 1$, $(\alpha)_n = \alpha(\alpha+1)\dots(\alpha+n-1)$, $4p = (q-1)^2$, $q \neq 1$, $\mathbf{z} = (z_1, z_2) \in \mathbb{C}^2$.

In the paper [4], formal expansions into branched continued fractions were constructed for the following functions:

$$\frac{H_4(a, b; c, b; \mathbf{z})}{H_4(a+1, b; c+1, b; \mathbf{z})} \quad (2)$$

and

$$\frac{H_4(a, d+1; c, d; \mathbf{z})}{H_4(a+1, d+1; c, d+1; \mathbf{z})}, \quad \frac{H_4(a, d+1; c, d; \mathbf{z})}{H_4(a, d+2; c, d+1; \mathbf{z})}. \quad (3)$$

In particular, it is proved that (2) has a formal expansion of the form

$$1 - z_2 - \frac{h_1 z_1}{1 - z_2 - \frac{h_2 z_1}{1 - z_2 - \frac{h_3 z_1}{1 - \dots}}}, \quad (4)$$

where

$$h_k = \frac{(2c - a + k - 1)(a + k)}{(c + k - 1)(c + k)}, \quad k \geq 1. \quad (5)$$

For $h_k, k \geq 1$, such that $0 < h_k \leq h$ for all $k \geq 1$, where h is a positive number, $c \notin \{0, -1, -2, \dots\}$, the convergence of branched continued fraction (4) to the function (2) was investigated in domain

$$H_h = \{z \in \mathbb{C}^2 : z_k \notin [1/(4 + 4h), +\infty), k = 1, 2\}.$$

In this paper, Theorem 2 provides convergence criteria for complex $h_k, k \geq 1$, while Theorems 3 and 4 for negative and positive $h_k, k \geq 1$, respectively. Since all the studied domains of convergence contain the neighborhood of the origin, the analytical extension of the functions to these domains is established by the PC method (see [1]). Section 2 provides several examples to illustrate this.

Some other recent studies related to the function (1) can be found in [14, 29, 32, 35].

1 Branched continued fraction and analytic continuation

In this section, we establish the domains of the analytic continuation of the function (1). Here we need the following theorem, proved in [15].

Theorem 1. *Let $m_{0,k}, k \geq 1$, be real numbers such that $0 < m_{0,k} \leq 1$ for all $k \geq 1$. Then the following is true.*

(A) *The branched continued fraction*

$$1 - z_{1,0} - \frac{m_{0,1} z_{0,1}}{1 - (1 - m_{0,1}) z_{1,1} - \frac{m_{0,2} (1 - m_{0,1}) z_{0,2}}{1 - (1 - m_{0,2}) z_{1,2} - \frac{m_{0,3} (1 - m_{0,2}) z_{0,3}}{1 - \dots}}} \quad (6)$$

converges for $|z_{1,k}| \leq 1/2$ and $|z_{0,k}| \leq 1/2$ for all $k \geq 1$.

(B) *If f_n denotes an n th approximant of (6), then $|f_n - 1| \leq 1$ for all $n \geq 1$.*

Note that various definitions of the convergence of branched continued fractions can be found in [15, 17].

Let us prove the following theorem.

Theorem 2. *Let a and c be complex constants such that*

$$|h_k| + \operatorname{Re}(h_k) \leq pq(1 - q) \quad \text{for all } k \geq 1, \quad (7)$$

where $h_k, k \geq 1$, are defined by (5), $c \notin \{0, -1, -2, \dots\}$, p is a positive number, and $0 < q < 1$. Then the following is true.

(A) The branched continued fraction (4) converges uniformly on every compact subset of the domain

$$H_{p,q}^{h,l} = H_{p,q} \cup H^{h,l}, \quad (8)$$

where

$$H_{p,q} = \left\{ \mathbf{z} \in \mathbb{C}^2 : |z_1| < \frac{1 + \cos(\arg(z_1))}{2p}, \operatorname{Re} \left(\frac{z_2}{e^{(i/2)\arg(z_1)}} \right) < \frac{q}{2} \cos \left(\frac{\arg(z_1)}{2} \right) \right\} \quad (9)$$

and

$$H^{h,l} = \left\{ \mathbf{z} \in \mathbb{C}^2 : |z_1| < \frac{l(1-l)}{2h}, |z_2| < \frac{1-l}{2} \right\}, \quad (10)$$

with

$$h = \max_{k \in \mathbb{N}} \{ |h_k| \}, \quad 0 < l < 1,$$

to a function $f(\mathbf{z})$ holomorphic in $H_{p,q}^{h,l}$.

(B) The function $f(\mathbf{z})$ is an analytic continuation of (2) in the domain (8).

Proof. An application of Theorem 1 (A), with $m_{0,k} = l$, $k \geq 1$, $0 < l < 1$, shows that the branched continued fraction (4) converges for all $\mathbf{z} \in H^{h,l}$. Theorem 1 (B) implies that the approximants of (4) all lie in $|w - 1| \leq 1$ if $\mathbf{z} \in H^{h,l}$. Hence, by [2, Theorem 3], the convergence of (4) is uniform on every compact subset of (10).

Let us prove the uniform convergence of the branched continued fraction (4) on every compact subset of (9) in the same way as in [18, Theorem 2].

We set $G_n^{(n)}(\mathbf{z}) = 1$, $n \geq 1$, and, for $1 \leq k \leq n-1$, $n \geq 2$,

$$G_k^{(n)}(\mathbf{z}) = 1 - z_2 - \frac{h_{k+1}z_1}{1 - z_2 - \frac{h_{k+2}z_1}{1 - \dots - z_2 - \frac{h_{n-1}z_1}{1 - z_2 - h_n z_1}}}.$$

It follows that

$$G_k^{(n)}(\mathbf{z}) = 1 - z_2 - \frac{h_{k+1}z_1}{G_{k+1}^{(n)}(\mathbf{z})}, \quad 1 \leq k \leq n-1, \quad n \geq 2, \quad (11)$$

and n th approximant of (4) we write as

$$f_n(\mathbf{z}) = 1 - z_2 - \frac{h_1 z_1}{G_1^{(n)}(\mathbf{z})}. \quad (12)$$

Let $\arg(z_1) = \varphi$, n be an arbitrary natural number, and \mathbf{z} be an arbitrary fixed point in the domain (9). Let us prove that

$$\operatorname{Re}(G_k^{(n)}(\mathbf{z})e^{-i\varphi/2}) > (1-q) \cos(\varphi/2) \geq c > 0, \quad 1 \leq k \leq n, \quad n \geq 1. \quad (13)$$

From an arbitrary fixed point \mathbf{z} , $\mathbf{z} \in H_{p,q}$, it follows that for its arbitrary neighborhood, there exists σ , $0 < \sigma \leq \pi/2$, such that $|\varphi/2| \leq \pi/2 - \sigma$ and, thus,

$$(1 - q) \cos(\varphi/2) \geq (1 - q) \cos(\varphi/2 - \sigma) = (1 - q) \sin(\sigma) = c > 0.$$

If $k = n$, the first inequality in (13) is obvious. Assuming that the first inequality in (13) is true if $k = s + 1 \leq n$. Then, for $k = s$ from (11) we obtain

$$G_s^{(n)}(\mathbf{z})e^{-i\varphi/2} = e^{-i\varphi/2} - z_2e^{-i\varphi/2} - \frac{h_{s+1}z_1e^{-i\varphi}}{G_{s+1}^{(n)}(\mathbf{z})e^{-i\varphi/2}}. \quad (14)$$

Using [1, Corollary 2], (7), (9), from (14) we get

$$\begin{aligned} \operatorname{Re}(G_s^{(n)}(\mathbf{z})e^{-i\varphi/2}) &\geq \cos(\varphi/2) - \operatorname{Re}(z_2e^{-i\varphi/2}) - \frac{|h_{s+1}| + \operatorname{Re}(h_{s+1})}{2 \operatorname{Re}(G_{s+1}^{(n)}(\mathbf{z})e^{-i\varphi/2})} |z_1| \\ &> \cos(\varphi/2) - \frac{q \cos(\varphi/2)}{2} - \frac{pq(1-q)}{2(1-q) \cos(\varphi/2)} \frac{1 + \cos(\varphi)}{2p} \\ &= (1 - q) \cos(\varphi/2). \end{aligned}$$

Thus, $G_1^{(n)}(\mathbf{z}) \neq 0$ for all $n \geq 1$ and $\mathbf{z} \in H_{p,q}$, i.e. that each approximant (12) is a holomorphic function in (9).

Now, let K be an arbitrary compact subset of (9), then there exists an open bi-disk

$$Q_R = \{\mathbf{z} \in \mathbb{C}^2 : |z_k| < R, k = 1, 2\}$$

of radius R , $R > 0$, such that $K \subset Q_R$. Then, for any $n \geq 1$ and $\mathbf{z} \in H_{p,q} \cap Q_R$ from (12) we get

$$|f_n(\mathbf{z})| \leq 1 + R + \frac{|h_1|R}{\operatorname{Re}(G_1^{(n)}(\mathbf{z})e^{-i\varphi/2})} < 1 + R + \frac{hR}{(1 - q) \cos(\varphi/2)} = C(K),$$

i.e. the sequence $\{f_n(\mathbf{z})\}$ is uniformly bounded on every compact subset of $H_{p,q}$.

It is easy to see that for every L such that

$$0 < L < \min \left\{ \frac{1-l}{2}, \frac{l(1-l)}{2h}, \frac{1}{p}, \frac{q}{2} \right\}$$

the domain

$$Y_L = \{\mathbf{z} \in \mathbb{R}^2 : 0 < z_k < L, k = 1, 2\}$$

contained in $H_{p,q}$, in particular $Y_{L/2} \subset H_{p,q}$. Using (7), for any $\mathbf{z} \in Y_L$, $Y_L \subset H_{p,q}$, it is easy to show that $|z_2| < (1 - l)/2$ and $|h_k z_1| < l(1 - l)/2$, $k \geq 1$, i.e. the elements of (4) satisfy Theorem 1, with $m_{0,k} = l$, $k \geq 1$. It shows that branched continued fraction (4) converges for all $\mathbf{z} \in Y_L$, $Y_L \subset H_{p,q}$. Hence, by [2, Theorem 3], the convergence of (4) is uniform on compact subsets of (9).

The proof of (B) is analogous to the proof of Theorem 2 (B) in [2], hence, it is omitted. \square

Remark 1. In Theorem 2, the set $\operatorname{Re}(z_2e^{-(i/2)\arg(z_1)}) < (q/2) \cos(\arg(z_1)/2)$ can also be written as $z_2 \notin [q/2, +\infty)$.

If we set $a = 0$ and replace c with $c - 1$ in Theorem 2, we get the following result.

Corollary 1. Let c be a complex constant satisfying (7), where

$$h_1 = \frac{2}{c}, \quad h_k = \frac{k(2c + k - 3)}{(c + k - 2)(c + k - 1)} \quad \text{for all } k \geq 2, \quad (15)$$

$c \notin \{0, -1, -2, \dots\}$, p is a positive number, and $0 < q < 1$. Then the branched continued fraction

$$\frac{1}{1 - z_2 - \frac{h_1 z_1}{1 - z_2 - \frac{h_2 z_1}{1 - \dots}}} \quad (16)$$

converges uniformly on every compact subset of (8) to a function $f(\mathbf{z})$ holomorphic in $H_{p,q}^{h,l}$, and $f(\mathbf{z})$ is an analytic continuation of $H_4(1, b; c, b; \mathbf{z})$ in the domain (8).

An application of Theorem 2 is the following theorem.

Theorem 3. Let a and c be real numbers such that $-h \leq h_k < 0$ for all $k \geq 1$, where $h_k, k \geq 1$, are defined by (5), h is a positive number. Then the branched continued fraction (4) converges uniformly on every compact subset of the domain

$$P_{h,l} = \left\{ \mathbf{z} \in \mathbb{C}^2 : z_1 \notin \left(-\infty, -\frac{l(1-l)}{2h} \right], z_2 \notin \left[\frac{1}{2}, +\infty \right) \right\}, \quad (17)$$

where $0 < l < 1$, to function $f(\mathbf{z})$ holomorphic in $P_{h,l}$, and $f(\mathbf{z})$ is an analytic continuation of (2) in the domain (17).

Proof. If $h_k < 0$ for all $k \geq 1$, then the condition (7) holds for all $p > 0$ and $0 < q < 1$. Let K be an arbitrary compact set contained in (17). Then $K \subseteq H_{p,q}^{h,l} \subseteq P_{h,l}$ for some p sufficiently small and q sufficiently close to 1, whose $H_{p,q}^{h,l}$ is the domain (8). Theorem 3 is, thus, an immediate consequence of Theorem 2. \square

Corollary 2. Let c be a real number such that $-h \leq h_k < 0$ for all $k \geq 1$, where $h_k, k \geq 1$, are defined by (15), h is a positive number. Then the branched continued fraction (16) converges uniformly on every compact subset of (17) to a function $f(\mathbf{z})$ holomorphic in $P_{h,l}$, and $f(\mathbf{z})$ is an analytic continuation of $H_4(1, b; c, b; \mathbf{z})$ in the domain (17).

The following result can be proved in much the same way as Theorem 2.

Theorem 4. Let a and c be real numbers such that $0 < h_k \leq h$ for all $k \geq 1$, where $h_k, k \geq 1$, are defined by (5), h is a positive number. Then the following is true.

(A) The branched continued fraction (4) converges uniformly on every compact subset of the domain

$$C_{h,l} = \left\{ \mathbf{z} \in \mathbb{C}^2 : z_1 \notin \left[\frac{1}{8h}, +\infty \right), z_2 \notin \left[\frac{1-l}{2}, +\infty \right) \right\}, \quad (18)$$

where $0 < l < 1$, to function $f(\mathbf{z})$ holomorphic in $C_{h,l}$.

(B) The function $f(\mathbf{z})$ is an analytic continuation of (2) in the domain (18).

Corollary 3. Let c be a real number such that $0 < h_k \leq h$ for all $k \geq 1$, where $h_k, k \geq 1$, are defined by (15), h is a positive number. Then the branched continued fraction (16) converges uniformly on every compact subset of (18) to a function $f(\mathbf{z})$ holomorphic in $C_{h,l}$, and $f(\mathbf{z})$ is an analytic continuation of $H_4(1, b; c, b; \mathbf{z})$ in the domain (18).

Remark 2. Results similar to Theorems 2–4 can be obtained for the expansions of the functions (3) constructed in paper [4].

2 Examples

As an example, by Corollary 3 we get

$$((1 - z_2)^2 - 4z_1)^{-1/2} = H_4(1, b; 1, b; \mathbf{z}) = \frac{1}{1 - z_2 - \frac{2z_1}{1 - z_2 - \frac{z_1}{1 - z_2 - \frac{z_1}{1 - \dots}}}}, \quad (19)$$

where the branched continued fraction converges and represents a single-valued branch of the analytic function on the left side of (19) in the domain (18) with $h = 2$.

One more example, by Corollary 3 we obtain

$$\begin{aligned} \arctan \frac{2\sqrt{-z_1}}{1 - z_2} &= 2\sqrt{-z_1}H_4(1, b; 3/2, b; \mathbf{z}) \\ &= \frac{2\sqrt{-z_1}}{1 - z_2 - \frac{\frac{4}{3}z_1}{1 - z_2 - \frac{\frac{16}{15}z_1}{1 - z_2 - \dots - \frac{\frac{4k^2}{4k^2 - 1}z_1}{1 - \dots}}}}, \end{aligned} \quad (20)$$

where the branched continued fraction converges and represents a single-valued branch of the analytic function of on the left side of (20) in the domain (18) with $h = 4/3$.

3 Conclusions

We consider the representation and extension of the analytic functions by special families of functions — branched continued fractions. The main results are new domains of analytical continuation for the Horn's hypergeometric function H_4 with certain conditions on real and complex parameters, which were established using their branched continued fraction representations. Further investigations can be continued in various directions. First, we can extend the domains of convergence of the branched continued fraction expansions with real and complex coefficients in their elements using parabolic domains of convergence [11, 12] and angular domains of convergence [9, 10]. Other research directions are the truncation errors analysis and the computational stability of the branched continued fraction expansions. Here, we can use the results of papers [7, 8, 13, 16, 21, 31].

References

- [1] Antonova T., Dmytryshyn R., Goran V. *On the analytic continuation of Lauricella-Saran hypergeometric function $F_K(a_1, a_2, b_1, b_2; a_1, b_2, c_3; \mathbf{z})$* . Mathematics 2023, **11** (21), 4487. doi:10.3390/math11214487
- [2] Antonova T., Dmytryshyn R., Kravtsiv V. *Branched continued fraction expansions of Horn's hypergeometric function H_3 ratios*. Mathematics 2021, **9** (2), 148. doi:10.3390/math9020148
- [3] Antonova T., Dmytryshyn R., Kril P., Sharyn S. *Representation of some ratios of Horn's hypergeometric functions H_7 by continued fractions*. Axioms 2023, **12** (8), 738. doi:10.3390/axioms12080738
- [4] Antonova T., Dmytryshyn R., Lutsiv I.-A., Sharyn S. *On some branched continued fraction expansions for Horn's hypergeometric function $H_4(a, b; c, d; z_1, z_2)$ ratios*. Axioms 2023, **12** (3), 299. doi:10.3390/axioms12030299

- [5] Antonova T., Dmytryshyn R., Sharyn S. *Branched continued fraction representations of ratios of Horn's confluent function H_6* . *Constr. Math. Anal.* 2023, **6** (1), 22–37. doi:10.33205/cma.1243021
- [6] Antonova T., Dmytryshyn R., Sharyn S. *Generalized hypergeometric function ${}_3F_2$ ratios and branched continued fraction expansions*. *Axioms* 2021, **10** (4), 310. doi:10.3390/axioms10040310
- [7] Antonova T.M., Dmytryshyn R.I. *Truncation error bounds for branched continued fraction whose partial denominators are equal to unity*. *Mat. Stud.* 2020, **54** (1), 3–14. doi:10.30970/ms.54.1.3-14
- [8] Bilanyk I.B. *A truncation error bound for some branched continued fractions of the special form*. *Mat. Stud.* 2019, **52** (2), 115–123. doi:10.30970/ms.52.2.115-123
- [9] Bodnar D.I., Bilanyk I.B. *Estimation of the rates of pointwise and uniform convergence of branched continued fractions with inequivalent variables*. *J. Math. Sci.* 2022, **265**, 423–437. doi:10.1007/s10958-022-06062-w (translation of *Mat. Metody Fiz.-Mekh. Polya* 2019, **62** (4), 72–82. (in Ukrainian))
- [10] Bodnar D.I., Bilanyk I.B. *On the convergence of branched continued fractions of a special form in angular domains*. *J. Math. Sci.* 2020, **246**, 188–200. doi:10.1007/s10958-020-04729-w (translation of *Mat. Metody Fiz.-Mekh. Polya* 2017, **63** (3), 60–69. (in Ukrainian))
- [11] Bodnar D.I., Bilanyk I.B. *Parabolic convergence regions of branched continued fractions of the special form*. *Carpathian Math. Publ.* 2021, **13** (3), 619–630. doi:10.15330/cmp.13.3.619-630
- [12] Bodnar D.I., Bilanyk I.B. *Two-dimensional generalization of the Thron-Jones theorem on the parabolic domains of convergence of continued fractions*. *Ukrainian Math. J.* 2023, **74** (9), 1317–1333. doi:10.1007/s11253-023-02138-1 (translation of *Ukrain. Mat. Zh.* 2022, **74** (9), 1155–1169. doi:10.37863/umzh.v74i9.7096 (in Ukrainian))
- [13] Bodnar D.I., Bodnar O.S., Bilanyk I.B. *A truncation error bound for branched continued fractions of the special form on subsets of angular domains*. *Carpathian Math. Publ.* 2023, **15** (2), 437–448. doi:10.15330/cmp.15.2.437-448
- [14] Blümlein J., Saragnese M., Schneider C. *Hypergeometric structures in Feynman integrals*. *Ann. Math. Artif. Intell.* 2023, **91**, 591–649. doi:10.1007/s10472-023-09831-8
- [15] Dmytryshyn R.I. *Convergence of multidimensional A- and J-fractions with independent variables*. *Comput. Methods Funct. Theory* 2022, **22** (2), 229–242. doi:10.1007/s40315-021-00377-6
- [16] Dmytryshyn R.I. *Convergence of some branched continued fractions with independent variables*. *Mat. Stud.* 2017, **47** (2), 150–159. doi:10.15330/ms.47.2.150-159
- [17] Dmytryshyn R., Goran V. *On the analytic extension of Lauricella-Saran's hypergeometric function F_K to symmetric domains*. *Symmetry* 2024, **16** (2), 220. doi:10.3390/sym16020220
- [18] Dmytryshyn R., Lutsiv I.-A., Bodnar O. *On the domains of convergence of the branched continued fraction expansion of ratio $H_4(a, d + 1; c, d; \mathbf{z}) / H_4(a, d + 2; c, d + 1; \mathbf{z})$* . *Res. Math.* 2023, **31** (2), 19–26. doi:10.15421/242311
- [19] Dmytryshyn R., Lutsiv I.-A., Dmytryshyn M., Cesarano C. *On some domains of convergence of branched continued fraction expansions of the ratios of Horn hypergeometric functions H_4* . *Ukr. Math. J.* 2024, **76** (4), 559–565. doi:10.1007/s11253-024-02338-3 (translation of *Ukrain. Mat. Zh.* 2024, **76** (4), 502–508. doi:10.3842/umzh.v74i4.7877 (in Ukrainian))
- [20] Dmytryshyn R.I., Lutsiv I.-A.V. *Three- and four-term recurrence relations for Horn's hypergeometric function H_4* . *Res. Math.* 2022, **30** (1), 21–29. doi:10.15421/242203
- [21] Dmytryshyn R.I. *On some of convergence domains of multidimensional S-fractions with independent variables*. *Carpathian Math. Publ.* 2019, **11** (1), 54–58. doi:10.15330/cmp.11.1.54-58
- [22] Dmytryshyn R.I. *On the expansion of some functions in a two-dimensional g-fraction with independent variables*. *J. Math. Sci.* 2012, **181** (3), 320–327. doi:10.1007/s10958-012-0687-5
- [23] Dmytryshyn R.I., Sharyn S.V. *Approximation of functions of several variables by multidimensional S-fractions with independent variables*. *Carpathian Math. Publ.* 2021, **13** (3), 592–607. doi:10.15330/cmp.13.3.592-607
- [24] Dmytryshyn R.I. *Two-dimensional generalization of the Rutishauser qd-algorithm*. *J. Math. Sci.* 2015, **208** (3), 301–309. doi:10.1007/s10958-015-2447-9

- [25] Horn J. *Hypergeometrische Funktionen zweier Veränderlichen*. Math. Ann. 1931, **105**, 381–407. doi:10.1007/BF01455825
- [26] Kaliuzhnyi-Verbovetskyi D., Pivovarchik V. *Recovering the shape of a quantum caterpillar tree by two spectra*. Mech. Math. Methods 2023, **5** (1), 14–24. doi:10.31650/2618-0650-2023-5-1-14-24
- [27] Kaminsky A.A., Selivanov M.F. *On the application of branched operator continued fractions for a boundary problem of linear viscoelasticity*. Int. Appl. Mech. 2006, **42**, 115–126. doi:10.1007/s10778-006-0066-3
- [28] Komatsu T. *Asymmetric circular graph with Hosoya index and negative continued fractions*. Carpathian Math. Publ. 2021, **13** (1), 608–618. doi:10.15330/cmp.13.3.608-618
- [29] Korkmaz-Duzgun D. *A new type multivariable multiple hypergeometric functions*. Turkish J. Math. Comput. Sci. 2021, **13** (2), 359–372. doi:10.47000/tjmcs.954676
- [30] Lima H. *Multiple orthogonal polynomials associated with branched continued fractions for ratios of hypergeometric series*. Adv. Appl. Math. 2023, **147**, 102505. doi:10.1016/j.aam.2023.102505
- [31] Manziy O., Hladun V., Ventyk L. *The algorithms of constructing the continued fractions for any ratios of the hypergeometric Gaussian functions*. Math. Model. Comput. 2017, **4** (1), 48–58. doi:10.23939/mmc2017.01.048
- [32] Parmar R.K., Choi J., Saravanan S. *Extended Exton's triple and Horn's double hypergeometric functions and associated bounding inequalities*. Symmetry 2023, **15** (6), 1132. doi:10.3390/sym15061132
- [33] Petreolle M., Sokal A.D. *Lattice paths and branched continued fractions II. Multivariate Lah polynomials and Lah symmetric functions*. Eur. J. Combin. 2021, **92**, 103235. doi:10.1016/j.ejc.2020.103235
- [34] Wang R., Qian J. *On branched continued fractions rational interpolation over pyramid-typed grids*. Numer. Algor. 2010, **54**, 47–72. doi:10.1007/s11075-009-9322-z
- [35] Younis J., Jain S., Agarwal P., Momani S. *Certain integral representations involving hypergeometric functions in two variables*. Math. Morav. 2022, **26** (1), 27–36. doi:10.5937/MatMor2201027Y

Received 21.12.2023

Revised 27.02.2024

Дмитришин Р., Луців І.-А., Дмитришин М. *Про аналітичне розширення гіпергеометричної функції Горна H_4 // Карпатські матем. публ. — 2024. — Т.16, №1. — С. 32–39.*

У статті досліджено нові області збіжності гіллястих ланцюгових дробових розвинень гіпергеометричної функції Горна H_4 з дійсними та комплексними параметрами. За допомогою РС методу встановлено аналітичні розширення деяких функцій у досліджені області збіжності. Для ілюстрації цього наприкінці наведено кілька прикладів.

Ключові слова і фрази: гіллястий ланцюговий дріб, гіпергеометрична функція Горна, апроксимація раціональними функціями, збіжність, аналітичне продовження.